

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/54, 15/82, 15/11, 9/10, A01H 5/00, C11B 1/00	A1	(11) International Publication Number: WO 94/13814 (43) International Publication Date: 23 June 1994 (23.06.94)
(21) International Application Number: PCT/GB93/02528 (22) International Filing Date: 10 December 1993 (10.12.93) (30) Priority Data: 9225845.8 10 December 1992 (10.12.92) GB (71) Applicant (for all designated States except US): NICKERSON BIOCEM LIMITED [GB/GB]; Cambridge Science Park, Milton Road, Cambridge CB4 4GZ (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): SLABAS, Antoni, Ryszard [GB/GB]; 8 Telford Close, High Shincliffe, Durham DH1 2YJ (GB). BROWN, Adrian, Paul [GB/GB]; 4 Church Villas, Shadforth, Durham DH6 1LQ (GB). (74) Agents: SHEARD, Andrew, Gregory et al.; Kilburn & Strode, 30 John Street, London WC1N 2DD (GB).		(81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: DNA ENCODING 2-ACYLTRANSFERASES (57) Abstract Plants, particularly transgenic plants, may be produced having a 2-acyltransferase enzyme or other insoluble acyltransferase enzyme with an altered substrate specificity compared to the native enzyme. For example, oil seed rape (<i>Brassica napus</i>) may contain a 2-acyltransferase transgene derived from <i>Limnanthes douglassi</i> in order to increase the erucic acid content of the oil. The cDNA sequence of maize (<i>Zea mays</i>) 2-acyltransferase is disclosed and is useful for cloning acyltransferase genes and/or cDNAs from other organisms, including <i>L. douglassi</i> .		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

DNA ENCODING 2-ACYLTRANSFERASES

This invention relates to modified plants. In particular, the invention relates to plants modified such that at least part of the plant (for example seeds of the plant) is capable of yielding a commercially useful oil.

Plants have long been a commercially valuable source of oil. Nutritional uses of plant-derived oils have hitherto been dominant, but attention is now turning additionally to plants as a source of industrially useful oils, for example as replacements for or improvements on mineral oils. Oil seeds, such as from rape, have a variety of lipids in them (Hildish & Williams, "Chemical Composition of Natural Lipids", Chapman Hall, London, 1964). There is now considerable interest in altering lipid composition by the use of recombinant DNA technology (Knauf, *TIBtech*, February 1987, 40-47), but by no means all of the goals have been realised to date for a variety of reasons, in spite of the ever-increasing sophistication of the technology.

Success in tailoring the lipid content of plant-derived oils requires a firm understanding of the biochemistry and genes involved. Broadly, two approaches are available. First, plants may be modified to permit the synthesis of fatty acids which are new (for the plant); so, for example, laurate and/or stearate may be synthesised in rape. Secondly, the pattern and/or extent of incorporation of fatty acids into the glycerol backbone of the lipid may be altered. It is with this latter approach that the present invention is concerned, although the former approach may additionally be used.

Lipids are formed in plants by the addition of fatty acid moieties onto the glycerol backbone by a series of acyl transferase enzymes. There are three positions on the glycerol molecule at which fatty acid (acyl) moieties may be substituted, and the substitution reached at each position is catalysed by a position-specific enzyme: the enzymes are known as 1-, 2- and 3-acyltransferases, respectively.

One, but not the only, current aim of "lipid engineering" in plants is to provide oils including lipids with a high content of erucic (22:1) acid. Erucic acid-containing lipids are commercially desirable for a number of purposes, particularly as replacements to or supplements for mineral oils in certain circumstances, as alluded to above. In the case of oil seed rape (*Brassica napus*), one of the most significant oil producing crops in cultivation today, the specificity of the 2-acyltransferase enzyme positively discriminates against the incorporation of erucic acid at position 2. So, even in those cultivars of rape which are able to incorporate erucic acid at positions 1 and 3, where there is no (or at least reduced) discrimination against erucic acid, only a maximum 66% of the fatty acids incorporated into triacyl glycerols can be erucic acid. Such varieties of rape are known as HEAR (high erucic acid rape) varieties.

It would therefore be desirable to increase the erucic acid content of conventional oil seed rape, as well as HEAR varieties; the same can be said of oils of other vegetable oil crops such as maize, sunflower and soya, to name but a few examples. While in principle it may be thought possible to introduce into a desired plant DNA encoding a 2-acyltransferase of different fatty acid

specificity, for example from a different plant, in practice there are a number of problems.

5 First, 2-acyltransferase and 3-acyltransferase are membrane bound, and therefore insoluble, enzymes. They have not been purified. This makes working with them difficult and rules out the use of many conventional DNA cloning procedures. This difficulty does not, paradoxically, lie in the way of cloning the gene (or at
10 least cDNA) encoding the 1-acyltransferase enzyme, which is soluble: in fact, recombinant DNA work has already been undertaken on this enzyme for a completely different purpose, namely the enhancement of chilling resistance in tobacco plant leaves, by Murata et al (Nature 356 710-713
15 (1992)).

Secondly, very little is known about the 2- and 3-acyltransferases. There is no idea of their size or how they are targeted to membranes. No nucleotide or amino
20 acid sequence data are available and no antibodies have been raised against them.

Although there has been discussion, therefore, of the desirability of modifying 2-acyltransferase specificity, for example by importing a gene coding for the
25 corresponding enzyme, but of different specificity, from another species, there is a pressing need in the art for the key which enables this work to be done. The present invention provides such a key, in the form of a DNA sequence (in the specific case, a cDNA sequence) encoding
30 a 2-acyltransferase. The DNA sequence in Figure 1 from nucleotides 130 to 1254 encodes the 2-acyltransferase from maize (*Zea mays*), including the stop codon.

According to a first aspect of the invention, therefore, there is provided a recombinant or isolated DNA sequence, preferably encoding an enzyme having membrane-bound acyltransferase activity, and selected from:

5

(i) a DNA sequence comprising the DNA sequence of Figure 1 encoding at least from MET₄₄ to Stop₄₁₈ (SEQ ID: 2) or its complementary strand,

10

(ii) nucleic acid sequences hybridising to the DNA sequence of Figure 1 (SEQ ID: 1), or its complementary strand, under stringent conditions, and

15

(iii) nucleic acid sequences which would hybridise to the DNA sequence of Figure 1 (SEQ ID: 1), or its complementary strand, but for the degeneracy of the genetic code.

20

Fragments of the above DNA sequences, for example of at least 15, 20, 30, 40 or 60 nucleotides in length, are also within the scope of the invention.

25

Suitable stringent conditions include salt solutions of approximately 0.9 molar at temperatures of from 35°C to 65°C. More particularly, stringent hybridisation conditions include 6 x SSC, 5 x Denhardt's solution, 0.5% SDS, 0.5% tetrasodium pyrophosphate and 50µg/ml denatured herring sperm DNA; washing may be for 2 x 30 minutes at 65°C in 1 x SSC, 0.1% SDS and 1 x 30 minutes in 0.2 x

30

SSC, 0.1% SDS at 65°C.

Nucleic acid sequences within the scope of the first aspect of the invention will generally encode a protein

having 2-acyltransferase activity, as that is the activity of the enzyme encoded by the Figure 1 nucleic acid sequence. Nucleic acid sequences not encoding a protein having enzymic activity (or the relevant enzymic activity) but otherwise conforming to the first aspect of the invention as set out above may be useful for other purposes (and are therefore also encompassed by the invention); for example they may be useful as probes, which is a utility shared by the nucleic acid sequences of the first aspect of the invention, including the Figure 1 sequence itself.

The probe utility arises as follows. As there is likely to be a high degree of homology between acyltransferases of different species (and particularly between 2-acyltransferases of different species) the Figure 1 sequence (or part of it, or other sequences within the invention) may be used to probe cDNA or genomic libraries of other species in order to clone DNA sequences encoding acyltransferases having desired specificities. For example, if it is desired to produce oil having a high content of erucic acid esterified to glycerol, a DNA library of any species which naturally makes erucic acid may be probed. Suitable plants include meadow foam (*Limnanthes* spp., especially *L. alba* and, particularly, *L. douglassi*) and *Crambe*. *Limnanthes douglassi* is the preferred species, as specificity studies show that there is positive discrimination towards incorporation of erucic acid into position 2 of the triacylglyceride. Libraries of organisms other than the higher plants may be probed; for example, certain bacteria may have an acyltransferase of the desired specificity.

DNA in accordance with the invention will in general have a higher degree of homology with at least part of the Figure 1 sequence than with known sequences.

5 Recombinant DNA in accordance with the invention may be in the form of a vector, which may have sufficient regulatory sequences (such as a promoter) to direct expression. Vectors which are not expression vectors are useful for cloning purposes (as expression vectors
10 themselves may be). Host cells (such as bacteria and plant cells) containing vectors in accordance with the invention themselves form part of the invention.

15 DNA sequences in accordance with the invention can be used in another way in cloning a gene of interest from another species: if the DNA is coupled to a suitable promoter, for example on an expression vector in a suitable host organism, protein may be produced. Such protein may be used to generate polyclonal or monoclonal
20 antibodies, or other binding molecules, which may then be used to screen for expression of homologous proteins in other species, for example as part of a DNA library screening programme.

25 Suitable cDNA libraries of target species will generally be prepared when the gene of interest is likely to be expressed; so cDNA embryo libraries (prepared at the early lipid synthesis stage), for example of *Limnanthes* spp. will be preferred.

30 The invention therefore enables the cloning of a wide variety of genes (or, more generally, DNA sequences) encoding acyltransferases, and 2-acyltransferases in particular, using DNA sequences as described above.

Such acyltransferases, such as from *Limnanthes* spp. may also be cloned directly, for example using complementation studies, from a DNA library of the species in question. For example, if *E. coli* is used as the complementation host, a mutant is chosen which is defective in the relevant enzyme (for example 2-acyltransferase); the DNA library from the target species (such as *L. douglassi*) is cloned into the mutant complementation host; host cells incorporating the target acyltransferase gene in their genome can readily be selected using appropriate selective media. *E. coli* mutant JC201 is a suitable host for use in complementation studies relating to 2-acyltransferase.

Cloning the acyltransferase gene of choice into a microbial host, such as a bacterium like *E. coli*, in such a way that the gene can be expressed has a particularly advantage in that the substrate specificity of the acyltransferase gene can be assessed in the microbial host before transformed plants are prepared, thereby saving considerably on research time. Such an assessment may be made by competitive substrate assays, in which differently detectably labelled candidate substrates for the enzyme compete with each other for incorporation into the glyceride. For example, ^{14}C -erucyl CoA and ^3H -oleoyl CoA can be used as competitive substrates for 2-acyltransferase, and the relative amounts of ^{14}C or tritium uptake into glyceride can be measured. (As 2-acyltransferases have acceptor, glycerol-based, substrates and donor, fatty acid-based, substrates, the experiment can be carried out with different acceptors, such as 1-erucyl-glycerol-3-phosphate and 1-oleoyl-glycerol-3-phosphate.) A gene coding for an enzyme which preferentially donates erucic acid to the acceptor

(particularly 1-erucyl-glycerol-3-phosphate) may by this means be identified as a DNA sequence of choice for further use in the invention as described below.

5 In a second aspect of the invention, there is provided a plant having one or more insoluble acyltransferase enzymes having a substrate specificity which differs from the native enzyme of the plant.

10 While site-directed mutagenesis and/or other protein engineering techniques may be used to alter the specificity of an enzyme native to the plant, it is preferred that the plant be transgenic and incorporate an expressible acyltransferase gene encoding an enzyme of
15 the desired specificity from another species. 2-acyltransferases are the enzymes of choice. For example, as described above, a 2-acyltransferase enzyme which has an enhanced specificity for, or at least no discrimination against, erucic acid, may be made by this
20 means to express in a plant which would not normally incorporate erucic acid into triacylglycerides. An important embodiment of the invention relates to genetically engineered plants which have higher levels of erucic acid incorporated into triacylglycerols than in
25 corresponding non-engineered plants. Preferable though this embodiment may be, though, the invention is not limited to the enhancement of erucic acid incorporation into glycerides: other acids may be desired in other circumstances.

30

For the acyltransferase transgene to be expressible, a promoter has to be operatively coupled to it. Because at the present state of the art it is difficult precisely to regulate the site of incorporation of a transgene into

the host genome, it is preferred that the transgene be coupled to its promoter prior to transformation of the plant. Promoters useful in the invention may be temporal- and/or seed-specific, but there is no need for them to be so: constitutive promoters, such as the CaMV 35S promoter, may in fact be preferred because they are usually strong promoters. Other tissues are unlikely to be adversely affected if the transgene encoding the acyltransferase enzyme is expressed in them, as the availability of the fatty acid CoA substrates is effectively limited to the seed.

The promoter-transgene construct, once prepared, is introduced into plant cells by any suitable means. The invention extends to such plant cells. Preferably, DNA is transformed into plant cells using a disarmed Ti-plasmid vector and carried by *Agrobacterium* by procedures known in the art, for example as described in EP-A-0116718 and EP-A-0270822. Alternatively, the foreign DNA could be introduced directly into plant cells using an electrical discharge apparatus. This method is preferred where *Agrobacterium* is ineffective, for example where the recipient plant is monocotyledonous. Any other method that provides for the stable incorporation of the DNA within the nuclear DNA of any plant cell of any species would also be suitable. This includes species of plant which are not currently capable of genetic transformation.

Preferably DNA in accordance with the invention also contains a second chimeric gene (a "marker" gene) that enables a transformed plant or tissue culture containing the foreign DNA to be easily distinguished from other plants or tissue culture that do not contain the foreign

DNA. Examples of such a marker gene include antibiotic resistance (Herrera-Estrella et al, *EMBO J.* 2(6) 987-95 (1983) and Herrera-Estrella et al, *Nature* 303 209-13 (1983)), herbicide resistance (EP-A-0242246) and glucuronidase (GUS) expression (EP-A-0344029). Expression of the marker gene is preferably controlled by a second promoter which allows expression in cells other than the tapetum, thus allowing selection of cells or tissue containing the marker at any stage of regeneration of the plant. The preferred second promoter is derived from the gene which encodes the 35S subunit of Cauliflower Mosaic Virus (CaMV) coat protein. However any other suitable second promoter could be used.

A whole plant can be regenerated from a single transformed plant cell, and the invention therefore provides transgenic plants (or parts of them, such as propagating material) including DNA in accordance with the invention as described above. The regeneration can proceed by known methods.

In one embodiment of the invention, the transgenic plant's native acyltransferase gene which corresponds to the transgene may be rendered at least partially inoperative or removed. So, if the transgene encodes a 2-acyltransferase, the plant's native 2-acyltransferase may be rendered inoperative by, for example, antisense or ribozyme techniques, as is known in the art.

By means of the invention, plants generating oil with a tailored lipid content may be produced. For example, the lipid composition of triacylglycerides in a plant may be substantially altered to produce triacylglycerides with a desired fatty acid (for example erucic acid) content

higher than has hitherto been possible. For example, oil seed rape (*B. napus*) may be transformed to produce oil whose triacylglyceride has an erucic acid content of over 70%.

5

It can readily be seen that plants with increased lipid levels may be produced by means of the invention. However, the invention is also useful for producing plants with decreased lipid levels, which may be desired if elevated protein and/or starch levels are required. Decreased lipid levels may be achieved by interfering with the proper functioning of a gene encoding a 2-acyltransferase, for example by antisense or ribozyme technology. (Such reduced-lipid plants may if desired be further engineered for higher protein and/or starch content, if wished.)

10

15

20

Promoters which naturally drive 2-acyltransferases may also be obtained by hybridisation and/or restriction and/or sequencing studies using the Figure 1 sequence.

25

30

The invention enables the production of protein encoded by DNA of the first aspect of the invention, should that be desired. The protein may be expressed by host cells harbouring DNA in the form of an expression vector. The protein, which may be an enzyme having 2-acyltransferase activity, may have an amino acid sequence which is identical to or homologous with the Figure 1 sequence. The degree of homology will generally be greater than that of known proteins, and may be at least 40, 50, 60, 70, 80, 90, 95 or 99%.

Preferred features of each aspect of the invention are as for each other aspect *mutatis mutandis*.

The invention is illustrated by the following examples. The examples refer to the accompanying drawings, in which:

5 FIGURE 1 shows the cDNA sequence derived in Example 1 (SEQ ID: 1) and its derived protein sequence (SEQ ID: 2).

10 FIGURE 2 shows a sequence alignment of part of the gene products of *plsB* (SEQ ID: 3) and *plsC* (SEQ ID: 4) with part of the sequence shown in Figure 1 (SEQ ID: 5), showing a conserved motif. *plsB* is the *E. coli* sn-glycerol-3-phosphate acyltransferase gene and *plsC* the 1-acyl-sn-glycerol-3-phosphate
15 acyltransferase gene of *E. coli*. Double points indicate exact matches between two sequences and a single point conservative amino acid substitutions. Stars indicate identical amino acids in all three sequences and residues conserved in two out of the
20 three sequences are marked by a + symbol.

25 FIGURES 3A, 3B and 3C: Membrane phospholipids from *E. coli* strains were extracted into chloroform and separated by 2-dimensional thin layer chromatography. The first dimension (ascending) was developed using chloroform:methanol:water (65:25:4) and the second dimension (left to right) developed with chloroform:methanol:acetic acid (65:25:10). Phospholipids were visualised by autoradiography for
30 16 hours at -70°C using Fuji RX film. The *E. coli* strains used were (Figure 3A): JC201 which carries a thermosensitive mutation in the 1-acyl-sn-glycerol-3-phosphate acyltransferase gene; (Figure 3B): JC201 containing the plasmid pPLSC, which

encodes the *E. coli* 1-acyl-sn-glycerol-3-phosphate acyltransferase gene; (Figure 3C): JC201 containing the plasmid whose cDNA insert sequence is shown in Figure 1. LPA, lysophosphatidic acid; PE, phosphatidylethanolamine; CL, cardiolipin; PG, phosphatidylglycerol; PA, phosphatidic acid; O, origin. 20% of the ^{32}P is incorporated in LPA in JC201 and all of the corresponding label is incorporated in PE in both of the other two strains.

FIGURE 4: Acyltransferase assays were performed using ^{32}P -labelled lysophosphatidic acid which had been extracted from the *E. coli* strain JC201 and oleoyl CoA as an acyl donor. Phospholipids present in the reaction mixtures were extracted into chloroform and separated using silica gel thin layer chromatography. Chloroform:methanol:acetic acid:water (25:15:4:2) was used to develop the plates. The phospholipids were visualised by autoradiography for 16 hours at -70°C using Fuji RX film. The *E. coli* strains used were: JC201 which carries a thermosensitive mutation in the 1-acyl-sn-glycerol-3-phosphate acyltransferase gene; JC201 containing the plasmid pPLSC which encodes the *E. coli* 1-acyl-sn-glycerol-3-phosphate acyltransferase gene; JC201 containing the plasmid whose maize cDNA insert sequence is shown in Figure 1. LPA, lysophosphatidic acid; PA, phosphatidic acid.

FIGURE 5 shows a comparison of the protein sequence shown in Figure 1 (SEQ ID: 6) with that derived (SEQ ID: 7) from a *B. napus* seed cDNA insert which was isolated by DNA hybridisation to the maize cDNA sequence. The sequences were aligned with the FastA

align program (1988). Double points signify identical amino acids and single points conservative amino acid substitutions.

5 Maize = 374 aa vs.rape = 311 aa
 51.5% identity; Optimised score: 705

EXAMPLE 1

Derivation of the Figure 1 cDNA sequence

10 Complementation studies using a maize cDNA expression library transferred into the *E. coli* mutant JC201 allowed the isolation of a plasmid encoding a 2-acyltransferase enzyme from maize. The cDNA insert of this plasmid is 1.6kb in size, and includes a poly A tail of 70bp. The
15 insert was sequenced to give the data shown and translation of the sequence revealed the presence of only one large open reading frame. This is shown on Figure 1 with proposed start methionine and stop codon boxed. The 2-acyltransferase is 374 amino acids in size and
20 sequencing upstream of open reading frame showed that the protein is expressed as part of a fusion protein in *E. coli*. This consists of 10 amino acids of the β -galactosidase protein, 43 amino acids (shown in sequence) corresponding to the 5' untranslated region of the mRNA and the 374 amino acid protein. Protein sequence
25 comparisons of the large open reading frame with the 2-acyltransferase of *E. coli* show little overall identity but there is a stretch of 80 residues which has a high level of conservative substitution and contains some
30 amino acids that are conserved in the 2-acyltransferase, 1-acyltransferase and N-acetyl glucosamine acyltransferase of *E. coli*.

EXAMPLE 2**Incorporation of ^{32}P into total phospholipids**

E. coli strains were grown in minimal medium containing ^{32}P orthophosphate. Total glycerolipids were extracted into organic solvents and separated by 2D thin layer chromatography (Figure 3) (Lysophosphatidic acid (LPA) is the substrate for 2-acyltransferase (2-AT)).

As can be seen in Figure 3A, the accumulation of ^{32}P -labelled LPA in the mutant JC 201 illustrates the absence of a fully functional 2-AT. Addition of a plasmid carrying either the native *E. coli* gene (Figure 3B), or the maize clone given in Figure 1 (Figure 3C) restores 2-AT activity to the cells, allowing LPA to be removed and further metabolised. (Lysophosphatidic acid.)

These data indicate that the DNA sequence given in Figure 1 codes for 2-AT.

EXAMPLE 3**Over expression of the cDNA**

The cDNA region specifying the protein sequence given in Figure 1 was cloned into the *E. coli* overexpression vector pET11d (Studier et al, *Meth. Enzymol.* 185 60-89 (1990)). Increased 2-acyltransferase activity following induction of expression from the plasmid insert confirmed that the sequence in Figure 1 is that of 2-AT.

EXAMPLE 4**Localisation of 2-AT activity in *E. coli* cells containing the maize clone**

2-acyltransferase assays were carried out using membranes isolated from the mutant strain JC.201 which lacks 2-AT and from JC.201 containing the maize plasmid (Figure 4).

2-AT activity was not detected in membrane fractions from JC.201. The addition of a plasmid carrying the native *E. coli* gene or the sequence given in Figure 1, to JC.201 resulted in restoration of 2-AT activity to the membranes.

EXAMPLE 5

Using the maize cDNA as a heterologous probe to obtain cDNA from oilseed rape

A seed cDNA library from *Brassica napus* was screened with the sequence given in Figure 1, using standard techniques (Sambrook et al "Molecular Cloning - A Laboratory Manual", 2nd Edition, Cold Spring Harbor Laboratory Press, 1989).

Conditions

For the hybridisation of the maize cDNA insert to the rape library: hybridisation was in 6xSSC, 5xDenhardt's solution, 0.5%SDS, 0.5% tetrasodium pyrophosphate and 50ugml⁻¹ denatured herring sperm DNA. The filters were washed 2 x 30 minutes at 65°C in 1xSSC, 0.1%SDS and 1 x 30 minutes in 0.2xSSC, 0.1%SDS at 65°C.

A hybridising clone was sequenced and a protein sequence derived for the large ORF. Alignment of this protein sequence with that derived from the maize cDNA clone given in Figure 1, is shown in Figure 5.

The strong identity between these sequences illustrates the potential of using the sequence given in Figure 1 to obtain other 2-ATS.

EXAMPLE 6**Transgenic plants**

The sequence given in Figure 1 can be cloned, alongside a suitable promoter, into a suitable vector for expression in plants. The vector can be used to transform plants and the resulting plants expressing the 2-AT can be analysed for lipid content. Lipid metabolism is expected to be upregulated and elevated lipid levels were detectable in seeds.

EXAMPLE 7**Antisense**

The sequence given in Figure 1 may be cloned, alongside a suitable promoter, in the antisense orientation into a suitable vector for expression in plants. The vector can be used to transform plants and the resulting plants expressing the 2-AT can be analysed for protein and starch content. Elevated levels of starch and protein are expected to be detectable in seeds.

EXAMPLE 8**Down-regulation of Native 2-AT**

The DNA sequence of a 2-AT derived from *L. douglassii* (obtained as described in Example 5) can be introduced into oilseed rape (OSR) under the expression of a suitable promoter, using vectors and plant transformation methods well known in the art. A second sequence, comprising antisense or ribozymes against the rape cDNA (Example 5) can be introduced for simultaneous expression. The resultant transformed plant is expected to have 2-AT activity corresponding to that of *L. douglassii*, with concurrent down regulation of the native rape 2-AT gene.

The modified OSR plant thus obtained had higher levels of erucic acid in position 2 of its triacylglycerols than wild type plants. In addition higher levels of trierucin are found in the seed oil.

5 -

EXAMPLE 9**Genomic library screening**

The sequence given in Figure 1 is used to screen a genomic library of *Arabidopsis* and a hybrid using clone
10 obtained. Using standard techniques, a promoter may be derived from this clone. The promoter may be used to drive expression in plant cell membranes.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

(A) NAME: Nickerson BIOCEM Limited
(non-US only)
(B) STREET: Cambridge Science Park, Milton Road
(C) CITY: Cambridge
(E) COUNTRY: United Kingdom
(F) POSTAL CODE (ZIP): CB4 4GZ

(A) NAME: SLABAS, Antoni Ryszard
(US only)
(B) STREET: 8 Telford Close, High Shinclffe
(C) CITY: Durham
(E) COUNTRY: United Kingdom
(F) POSTAL CODE (ZIP): DH1 2YJ

(A) NAME: BROWN, Adrian Paul
(US only)
(B) STREET: 4 Church Villas, Shadforth
(C) CITY: Durham
(E) COUNTRY: United Kingdom
(F) POSTAL CODE (ZIP): DH6 1LQ

(ii) TITLE OF INVENTION: DNA ENCODING 2-ACYLTRANSFERASES

(iii) NUMBER OF SEQUENCES: 7

(iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25 (EPO)

(v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: WO PCT/GB93/_____

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1514 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 130..1254

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CCCCGTCCTC CTCGTCGCCG GCGGAGCCGC CTACTATCGC CTGGAGAAGG AGCGCCGCGG	60
GGAGCTTTTC CCACTGCCGA CTGCCGTCTG ACCCTCCGAG ATCGGAAGCG GCGCCGGCGC	120
CGGCCGGCG ATG GCG ATC CCG CTC GTG CTC GTC GTG CTC CCG CTC GGC	168
Met Ala Ile Pro Leu Val Leu Val Val Leu Pro Leu Gly	
1 5 10	

20

CTG	CTC	TTC	CTC	CTG	TCC	GGC	CTC	ATC	GTC	AAC	GCC	ATC	CAG	GCC	GTC	216
Leu	Leu	Phe	Leu	Leu	Ser	Gly	Leu	Ile	Val	Asn	Ala	Ile	Gln	Ala	Val	
15						20					25					
CTA	TTT	GTG	ACG	ATA	AGG	CCC	TTT	TCG	AAG	AGC	TTC	TAC	CGT	CGG	ATC	264
Leu	Phe	Val	Thr	Ile	Arg	Pro	Phe	Ser	Lys	Ser	Phe	Tyr	Arg	Arg	Ile	
30					35				40						45	
AAC	AGA	TTC	TTG	GCC	GAG	CTG	CTG	TGG	CTT	CAG	CTT	GTC	TGG	GTG	GTG	312
Asn	Arg	Phe	Leu	Ala	Glu	Leu	Leu	Trp	Leu	Gln	Leu	Val	Trp	Val	Val	
				50					55					60		
GAC	TGG	TGG	GCA	GGT	GTT	AAG	GTA	CAA	CTG	CAT	GCA	GAT	GAG	GAA	ACT	360
Asp	Trp	Trp	Ala	Gly	Val	Lys	Val	Gln	Leu	His	Ala	Asp	Glu	Glu	Thr	
			65					70					75			
TAC	AGA	TCA	ATG	GGT	AAA	GAG	CAT	GCA	CTC	ATC	ATA	TCA	AAT	CAT	CGG	408
Tyr	Arg	Ser	Met	Gly	Lys	Glu	His	Ala	Leu	Ile	Ile	Ser	Asn	His	Arg	
	80						85					90				
AGT	GAT	ATT	GAT	TGG	CTC	ATT	GGA	TGG	ATA	TTG	GCC	CAG	CGT	TCA	GGG	456
Ser	Asp	Ile	Asp	Trp	Leu	Ile	Gly	Trp	Ile	Leu	Ala	Gln	Arg	Ser	Gly	
95						100					105					
TGC	CTT	GGA	AGT	ACA	CTT	GCT	GTC	ATG	AAG	AAG	TCA	TCC	AAG	TTC	CTT	504
Cys	Leu	Gly	Ser	Thr	Leu	Ala	Val	Met	Lys	Lys	Ser	Ser	Lys	Phe	Leu	
110					115					120					125	
CCA	GTT	ATT	GGC	TGG	TCA	ATG	TGG	TTT	GCA	GAG	TAC	CTC	TTT	TTG	GAA	552
Pro	Val	Ile	Gly	Trp	Ser	Met	Trp	Phe	Ala	Glu	Tyr	Leu	Phe	Leu	Glu	
				130					135					140		
AGG	AGC	TGG	GCC	AAG	GAT	GAA	AAG	ACA	CTA	AAG	TGG	GGT	CTC	CAA	AGG	600
Arg	Ser	Trp	Ala	Lys	Asp	Glu	Lys	Thr	Leu	Lys	Trp	Gly	Leu	Gln	Arg	
			145					150					155			
TTG	AAA	GAC	TTC	CCT	AGA	CCA	TTT	TGG	CTA	GCT	CTT	TTC	GTC	GAG	GGT	648
Leu	Lys	Asp	Phe	Pro	Arg	Pro	Phe	Trp	Leu	Ala	Leu	Phe	Val	Glu	Gly	
	160						165					170				
ACT	CGC	TTT	ACT	CCA	GCA	AAG	CTT	CTC	GCA	GCT	CAG	GAA	TAT	GCG	GCC	696
Thr	Arg	Phe	Thr	Pro	Ala	Lys	Leu	Leu	Ala	Ala	Gln	Glu	Tyr	Ala	Ala	
	175					180					185					
TCC	CAG	GGC	TTA	CCG	GCT	CCT	AGA	AAT	GTA	CTT	ATT	CCA	CGT	ACC	AAG	744
Ser	Gln	Gly	Leu	Pro	Ala	Pro	Arg	Asn	Val	Leu	Ile	Pro	Arg	Thr	Lys	
190					195				200						205	
GGA	TTT	GTA	TCT	GCT	GTA	AGT	ATT	ATG	CGA	GAT	TTT	GTT	CCA	GCC	ATT	792
Gly	Phe	Val	Ser	Ala	Val	Ser	Ile	Met	Arg	Asp	Phe	Val	Pro	Ala	Ile	
				210					215					220		
TAT	GAT	ACA	ACT	GTA	ATA	GTC	CCT	AAA	GAT	TCC	CCT	CAA	CCA	ACA	ATG	840
Tyr	Asp	Thr	Thr	Val	Ile	Val	Pro	Lys	Asp	Ser	Pro	Gln	Pro	Thr	Met	
			225				230						235			
CTG	CGG	ATT	TTG	AAA	GGG	CAA	TCA	TCA	GTG	ATA	CAT	GTC	CGC	ATG	AAA	888
Leu	Arg	Ile	Leu	Lys	Gly	Gln	Ser	Ser	Val	Ile	His	Val	Arg	Met	Lys	
		240					245					250				
CGT	CAT	GCA	ATG	AGT	GAG	ATG	CCA	AAA	TCA	GAT	GAG	GAT	GTT	TCA	AAA	936
Arg	His	Ala	Met	Ser	Glu	Met	Pro	Lys	Ser	Asp	Glu	Asp	Val	Ser	Lys	
	255					260					265					

TGG TGT AAA GAC ATT TTT GTG GCA AAG GAT GCC TTA CTG GAC AAG CAT	984
Trp Cys Lys Asp Ile Phe Val Ala Lys Asp Ala Leu Leu Asp Lys His	
270 275 280 285	
TTG GCA ACA GGC ACT TTC GAT GAG GAG ATT AGA CCT ATT GGC CGT CCA	1032
Leu Ala Thr Gly Thr Phe Asp Glu Glu Ile Arg Pro Ile Gly Arg Pro	
290 295 300	
GTG AAA TCA TTG CTG GTG ACC CTG TTC TGG TCG TGC CTC CTG CTG TTT	1080
Val Lys Ser Leu Leu Val Thr Leu Phe Trp Ser Cys Leu Leu Leu Phe	
305 310 315	
GGC GCC ATC GAG TTC TTC AAG TGG ACA CAG CTT CTG TCG ACG TGG AGG	1128
Gly Ala Ile Glu Phe Phe Lys Trp Thr Gln Leu Leu Ser Thr Trp Arg	
320 325 330	
GGT GTG GCG TTC ACT GCC GCA GGG ATG GCG CTT GTG ACG GGT GTC ATG	1176
Gly Val Ala Phe Thr Ala Ala Gly Met Ala Leu Val Thr Gly Val Met	
335 340 345	
CAT GTC TTC ATC ATG TTC TCC CAG GCT GAG CGG TCG AGC TCA GCC AGG	1224
His Val Phe Ile Met Phe Ser Gln Ala Glu Arg Ser Ser Ser Ala Arg	
350 355 360 365	
GCG GCA CGG AAC CGG GTC AAG AAG GAA TGAAAAATGG AGGGTGGAGA	1271
Ala Ala Arg Asn Arg Val Lys Lys Glu	
370 375	
TGAGGTTCTC GTGGGGTTTG TTATGGGCAA CCTTCAAAAG GACTCTCCAT TCATATTAGT	1331
ATTAATTCAT ATATATGCAG CGCCAAATTC CAGACATTGA TATGCTCTCA AATAGGATGT	1391
TCTGCTCCCC TCTTGATATT GTATGCAGGA AAGGGTTTGT AGGGAGTTTA CCCCCCCCCC	1451
CCCCCCCCCC GCCTTTCTTT GGGGAAGAAA GACATATTCT GGAAGCCTTC CAGTAGTTCA	1511
AAA	1514

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 374 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

```

Met Ala Ile Pro Leu Val Leu Val Val Leu Pro Leu Gly Leu Leu Phe
 1           5           10           15
Leu Leu Ser Gly Leu Ile Val Asn Ala Ile Gln Ala Val Leu Phe Val
 20           25           30
Thr Ile Arg Pro Phe Ser Lys Ser Phe Tyr Arg Arg Ile Asn Arg Phe
 35           40           45
Leu Ala Glu Leu Leu Trp Leu Gln Leu Val Trp Val Val Asp Trp Trp
 50           55           60
Ala Gly Val Lys Val Gln Leu His Ala Asp Glu Glu Thr Tyr Arg Ser
 65           70           75           80
Met Gly Lys Glu His Ala Leu Ile Ile Ser Asn His Arg Ser Asp Ile
 85           90           95
Asp Trp Leu Ile Gly Trp Ile Leu Ala Gln Arg Ser Gly Cys Leu Gly
 100          105          110
Ser Thr Leu Ala Val Met Lys Lys Ser Ser Lys Phe Leu Pro Val Ile
 115          120          125
Gly Trp Ser Met Trp Phe Ala Glu Tyr Leu Phe Leu Glu Arg Ser Trp
 130          135          140
Ala Lys Asp Glu Lys Thr Leu Lys Trp Gly Leu Gln Arg Leu Lys Asp
 145          150          155          160
Phe Pro Arg Pro Phe Trp Leu Ala Leu Phe Val Glu Gly Thr Arg Phe
 165          170          175
Thr Pro Ala Lys Leu Leu Ala Ala Gln Glu Tyr Ala Ala Ser Gln Gly
 180          185          190
Leu Pro Ala Pro Arg Asn Val Leu Ile Pro Arg Thr Lys Gly Phe Val
 195          200          205
Ser Ala Val Ser Ile Met Arg Asp Phe Val Pro Ala Ile Tyr Asp Thr
 210          215          220
Thr Val Ile Val Pro Lys Asp Ser Pro Gln Pro Thr Met Leu Arg Ile
 225          230          235          240
Leu Lys Gly Gln Ser Ser Val Ile His Val Arg Met Lys Arg His Ala
 245          250          255
Met Ser Glu Met Pro Lys Ser Asp Glu Asp Val Ser Lys Trp Cys Lys
 260          265          270
Asp Ile Phe Val Ala Lys Asp Ala Leu Leu Asp Lys His Leu Ala Thr
 275          280          285

```

23

Gly Thr Phe Asp Glu Glu Ile Arg Pro Ile Gly Arg Pro Val Lys Ser
 290 295 300
 Leu Leu Val Thr Leu Phe Trp Ser Cys Leu Leu Leu Phe Gly Ala Ile
 305 310 315 320
 Glu Phe Phe Lys Trp Thr Gln Leu Leu Ser Thr Trp Arg Gly Val Ala
 325 330 335
 Phe Thr Ala Ala Gly Met Ala Leu Val Thr Gly Val Met His Val Phe
 340 345 350
 Ile Met Phe Ser Gln Ala Glu Arg Ser Ser Ser Ala Arg Ala Ala Arg
 355 360 365
 Asn Arg Val Lys Lys Glu
 370

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Tyr	Phe	Val	Glu	Gly	Gly	Arg	Ser	Arg	Thr	Gly	Arg	Leu	Leu	Asp
1				5					10					15

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met	Phe	Pro	Glu	Gly	Thr	Arg	Ser	Arg	Gly	Arg	Gly	Leu	Leu	Pro
1				5					10					15

(2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

```

Leu Phe Val Glu Gly Thr Arg Phe Thr Pro Ala Lys Leu Leu Ala
1           5           10           15

```

(2) INFORMATION FOR SEQ ID NO: 6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 374 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

```

Met Ala Ile Pro Leu Val Leu Val Val Leu Pro Leu Gly Leu Leu Phe
1           5           10           15

Leu Leu Ser Gly Leu Ile Val Asn Ala Ile Gln Ala Val Leu Phe Val
20          25          30

Thr Ile Arg Pro Phe Ser Lys Ser Phe Tyr Arg Arg Ile Asn Arg Phe
35          40          45

Leu Ala Glu Leu Leu Trp Leu Gln Leu Val Trp Val Val Asp Trp Trp
50          55          60

Ala Gly Val Lys Val Gln Leu His Ala Asp Glu Glu Thr Tyr Arg Ser
65          70          75          80

Met Gly Lys Glu His Ala Leu Ile Ile Ser Asn His Arg Ser Asp Ile
85          90          95

Asp Trp Leu Ile Gly Trp Ile Leu Ala Gln Arg Ser Gly Cys Leu Gly
100         105         110

Ser Thr Leu Ala Val Met Lys Lys Ser Ser Lys Phe Leu Pro Val Ile
115        120        125

Gly Trp Ser Met Trp Phe Ala Glu Tyr Leu Phe Leu Glu Arg Ser Trp
130        135        140

Ala Lys Asp Glu Lys Thr Leu Lys Trp Gly Leu Gln Arg Leu Lys Asp
145        150        155        160

Phe Pro Arg Pro Phe Trp Leu Ala Leu Phe Val Glu Gly Thr Arg Phe
165        170        175

Thr Pro Ala Lys Leu Leu Ala Ala Gln Glu Tyr Ala Ala Ser Gln Gly
180        185        190

```

27

Leu Pro Ala Pro Arg Asn Val Leu Ile Pro Arg Thr Lys Gly Phe Val
 195 200 205
 Ser Ala Val Ser Ile Met Arg Asp Phe Val Pro Ala Ile Tyr Asp Thr
 210 215 220
 Thr Val Ile Val Pro Lys Asp Ser Pro Gln Pro Thr Met Leu Arg Ile
 225 230 235 240
 Leu Lys Gly Gln Ser Ser Val Ile His Val Arg Met Lys Arg His Ala
 245 250 255
 Met Ser Glu Met Pro Lys Ser Asp Glu Asp Val Ser Lys Trp Cys Lys
 260 265 270
 Asp Ile Phe Val Ala Lys Asp Ala Leu Leu Asp Lys His Leu Ala Thr
 275 280 285
 Gly Thr Phe Asp Glu Glu Ile Arg Pro Ile Gly Arg Pro Val Lys Ser
 290 295 300
 Leu Leu Val Thr Leu Phe Trp Ser Cys Leu Leu Leu Phe Gly Ala Ile
 305 310 315 320
 Glu Phe Phe Lys Trp Thr Gln Leu Leu Ser Thr Trp Arg Gly Val Ala
 325 330 335
 Phe Thr Ala Ala Gly Met Ala Leu Val Thr Gly Val Met His Val Phe
 340 345 350
 Ile Met Phe Ser Gln Ala Glu Arg Ser Ser Ser Ala Arg Ala Ala Arg
 355 360 365
 Asn Arg Val Lys Lys Glu
 370

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 295 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

```

Met Ala Met Ala Ala Val Ile Val Pro Leu Gly Ile Leu Phe Phe
1           5           10          15
Ile Ser Gly Leu Val Val Asn Leu Leu Gln Arg Ser Gly Cys Leu Gly
20          25          30
Ser Ala Leu Ala Val Met Lys Lys Ser Ser Lys Phe Leu Pro Val Ile
35          40          45
Gly Trp Ser Met Trp Phe Ser Glu Tyr Leu Phe Leu Glu Arg Asn Trp
50          55          60
Ala Lys Asp Glu Ser Thr Leu Lys Ser Gly Leu Gln Arg Leu Asn Asp
65          70          75          80
Phe Pro Arg Pro Phe Trp Leu Ala Leu Phe Val Glu Gly Thr Arg Phe
85          90          95
Thr Glu Ala Lys Leu Lys Ala Ala Gln Glu Tyr Ala Ala Ser Ser Glu
100         105         110
Leu Pro Val Pro Arg Asn Val Leu Ile Pro Arg Thr Lys Gly Phe Val
115        120        125
Ser Ala Val Ser Asn Met Arg Ser Phe Val Pro Ala Ile Tyr Asp Met
130        135        140
Thr Val Ala Ile Pro Lys Thr Ser Pro Pro Pro Thr Met Leu Arg Leu
145        150        155        160
Phe Lys Gly Gln Pro Ser Val Val His Val His Ile Lys Cys His Ser
165        170        175
Met Lys Asp Leu Pro Glu Ser Glu Asp Glu Ile Ala Gln Trp Cys Arg
180        185        190
Asp Gln Phe Val Thr Lys Asp Ala Leu Leu Asp Lys His Ile Ala Ala
195        200        205
Asp Thr Phe Ala Gly Gln Lys Glu Gln Asn Ile Gly Arg Pro Ile Lys
210        215        220
Ser Leu Ala Val Val Leu Ser Trp Ala Cys Leu Leu Thr Leu Gly Ala
225        230        235        240
Met Lys Phe Leu His Trp Ser Asn Leu Phe Ser Ser Trp Lys Gly Ile
245        250        255
Ala Leu Ser Ala Leu Gly Leu Gly Ile Ile Thr Leu Cys Met Gln Ile
260        265        270
Leu Ile Arg Ser Ser Gln Ser Glu Arg Ser Thr Pro Ala Lys Val Ala
275        280        285

```

Pro Ala Lys Pro Lys Asp Asn
290 295

CLAIMS

1. A recombinant or isolated DNA sequence selected from
 - (i) a DNA sequence comprising the DNA sequence of Figure 1 encoding at least from Met₄₄ to Stop₄₁₈ (SEQ ID: 2) or its complementary strand;
 - (ii) nucleic acid sequences hybridising to the DNA sequence of Figure 1 (SEQ ID: 1) or its complementary strand under stringent conditions; and
 - (iii) nucleic acid sequences which would hybridise to the DNA sequence of Figure 1 (SEQ ID: 1), or its complementary strand, but for the degeneracy of the genetic code.
2. A DNA sequence as claimed in claim 1 which encodes an enzyme having membrane-bound acyltransferase activity.
3. A DNA sequence as claimed in claim 2, in which the acyltransferase activity is 2-acyltransferase activity.
4. An isolated protein which is the expression product of a DNA sequence as claimed in claim 1, 2 or 3.
5. An antibody capable of specifically binding to a protein as claimed in claim 4.
6. A plant having one or more insoluble acyltransferase enzymes having a substrate specificity which differs from the native enzyme of the plant.

7. A plant as claimed in claim 6, which is transgenic for an insoluble acyltransferase enzyme from another species.

5 8. A plant as claimed in claim 6 or 7, which is rape, maize, sunflower or soya.

10 9. A plant as claimed in claim 7, wherein the said other species is a species of the genus *Limnanthes* (such as *L. alba* or, preferably, *L. douglassi*) or *Crambe*.

15 10. A plant as claimed in any one of claims 6 to 9, wherein the insoluble acyltransferase enzyme is 2-acyltransferase.

11. A plant as claimed in claim 10, wherein the 2-acyltransferase has a higher specificity for erucic acid than the native enzyme of the plant.

20 12. A plant as claimed in any one of claims 6 to 11, wherein the said native enzyme is at least partially rendered inoperative or removed, for example by a ribozyme or by antisense nucleic acid.

25 13. A method of generating oil, the method comprising cultivating a plant as claimed in anyone of claims 6 to 12 and harvesting oil produced by the plant or a part (particularly seeds) thereof.

30 14. A microbial host capable of expressing a DNA sequence as claimed in claim 1, 2 or 3.

15. A protein which is substantially homologous to a protein as claimed in claim 4.

16. A fragment of a DNA sequence as claimed in claim 1, 2 or 3, comprising at least 15 nucleotides.

5 17. DNA encoding RNA which is antisense to RNA encoded by DNA as claimed in claim 1, 2 or 3.

18. DNA encoding a ribozyme specific to RNA encoded by DNA as claimed in claim 1, 2 or 3.

10 19. Isolated or recombinant DNA containing a promoter which naturally drives expression of a gene to produce a protein as claimed in claim 4 or 15.

1/8

FIG. 1 (1/3)

1/1	CCC CGT	CCT CCT	CGT CGC	CGG CGG	AGC CGC	CTA CTA	TCG CCT	GGA GAA	GGA GCG	CGG CCG	CGG CGG
	P R	P P	R R	R R	S R	L L	S P	G E	G A	P R	
31/11											
61/21	GGA GCT	TTT CCC	ACT GCC	GAC TGC	CGT CTG	ACC CTC	CGA GAT	CGG AAG	CGG CGG	CGG CGG	CGG CGC
	G A	F P	T A	D C	R L	T L	R D	R K	R R	R R	R R
91/31											
121/41	CGG CCG	GCG ATG	GCG ATC	CCG CTC	GTG CTC	GTG GTG	CTC CCG	CTC GGC	CTG CTG	CTC TTC	CTC CTC
	R P	A M	A I	P L	V L	V V	L P	L G	L L	L F	L L
151/51											
181/61	CTG TCC	GGC CTC	ATC GTC	AAC GCC	ATC CAG	GCC GTC	CTA TTT	GTG AGC	ATA AGG	CCC CCC	TTT F
	L S	G L	I V	N A	I Q	A V	L F	V T	I I	R P	F F
271/91											
241/81	TCG AAG	AGC TTC	TAC CGT	CGG ATC	AAC AGA	TTC TTG	GCC GAG	CTG CTG	TGG TGG	CTT CAG	CTT L
	S K	S F	Y R	R I	N R	F L	A E	L L	W L	Q L	L L
331/111											
301/101	GTC TGG	GTG GTG	GAC TGG	TGG GCA	GGT GTT	AAG GTA	CAA CTG	CAT GCA	GAT GAT	GAG GAA	ACT T
	V W	V V	D W	W A	G V	K V	Q L	H A	D E	E E	T T
391/131											
361/121	TAC AGA	ATG GGT	AAA GAG	CAT GCA	CTC CTC	ATC ATA	TCA AAT	CAT CSG	AGT AGT	GAT ATT	GAT D
	Y R	S M	G K	E H	A L	I I	S N	H R	S S	D I	D D
451/151											
421/141	TGG CTC	ATT GGA	TGG ATA	TTG GCC	CAG CGT	TCA GGG	TGC CTT	GGA AGT	ACA ACA	CTT GCT	GTC V
	W L	I G	W I	L A	Q R	S G	C L	G S	T T	L A	V V
511/171											
481/161	ATG AAG	AAG TCA	TCC AAG	TTC CTT	CCA GTT	ATT GGC	TGG TCA	ATG TGG	TTT TTT	GCA GAG	TAC Y
	M K	K S	S K	F L	P V	I G	W S	M W	F F	A A	E Y
571/191											
541/181	CTC TTT	TTG GAA	AGG AGC	TGG GCC	AAG GAT	GAA AAG	ACA CTA	AAG TGG	GGT GGT	CTC CAA	AGG R
	L F	L E	R S	W A	K D	E K	T L	K W	G G	L Q	R R

2/8

FIG.1
(2/3)

601/201	TTG AAA	GAC	TTT	CCT	AGA	CCA	TTT	TGG	CTA	GCT	CTT	631/211	TTC	GTC	GAG	GGT	ACT	CGC	TTT	ACT
L K	D	F	P	P	R	P	F	W	L	A	L		F	V	E	G	T	R	F	T
661/221	CCA GCA	AAG	CTT	CTC	GCA	GCT	CAG	GAA	TAT	GCG	GCC	691/231	TCC	CAG	GGC	TTA	CCG	GCT	CCT	AGA
P A	K	L	L	A	A	Q	E	Y	A	A	A		S	Q	G	L	P	A	P	R
721/241	AAT GTA	CTT	ATT	CCA	CGT	ACC	AAG	GGA	TTT	GTA	TCT	751/251	GCT	GTA	AGT	ATT	ATG	CGA	GAT	TTT
N V	L	I	P	R	T	K	G	F	V	S	A		A	V	S	I	M	R	D	F
781/261	GTT CCA	GCC	ATT	TAT	GAT	ACA	ACT	GTA	ATA	GTC	CCT	811/271	AAA	GAT	TCC	CCT	CAA	CCA	ACA	ATG
V P	A	I	Y	D	T	T	V	I	V	P	K		K	D	S	P	Q	P	T	M
841/281	CTG CGG	ATT	TTG	AAA	GGG	CAA	TCA	TCA	GTG	ATA	CAT	871/291	GTC	CGC	ATG	AAA	CGT	CAT	GCA	ATG
L R	I	L	K	G	Q	S	S	V	I	H	V		V	R	M	K	R	H	A	M
901/301	AGT GAG	ATG	CCA	AAA	TCA	GAT	GAG	GAT	GTT	TCA	AAA	931/311	TGG	TGT	AAA	GAC	ATT	TTT	GTG	GCA
S E	M	P	K	S	D	E	D	V	S	K	W		C	K	D	I	F	V	A	
961/321	AAG GAT	GCC	TTA	CTG	GAC	AAG	CAT	TTG	GCA	ACA	GGC	991/331	ACT	TTC	GAT	GAG	GAG	ATT	AGA	CCT
K D	A	L	L	L	D	K	H	L	A	T	G		T	F	D	E	E	I	R	P
1021/341	ATT GGC	CGT	CCA	GTG	AAA	TCA	TTG	CTG	GTG	ACC	CTG	1051/351	TTC	TGG	TCG	TGC	CTC	CTG	CTG	TTT
I G	R	P	V	K	S	L	L	V	T	L	F		F	W	S	C	L	L	L	F
1081/361	GGC GCC	ATC	GAG	TTT	TTC	AAG	TGG	ACA	CAG	CTT	CTG	1111/371	TCG	ACG	TGG	AGG	GGT	GTG	GCG	TTC
G A	I	E	F	F	K	W	T	Q	L	L	S		T	W	R	G	V	A	F	

SUBSTITUTE SHEET

3/8




1141/381
 ACT GCC GCA GGG ATG GCG CTT GTG ACG GGT GTC ATG CAT GTC TTC ATC ATG TTC TCC CAG
 T A A G M A L A T V G G V M H V F I M F S Q
 1201/401
 GCT GAG CGG TCG AGC TCA GCC AGG GCG GCA CGG AAC CGG GTC AAG AAG GAA TGA AAA ATG
 A E R S S S A R A A R N R V K K E * K M
 1261/421
 GAG GGT GGA GAT GAG GTT CTC GTG GGG TTT GTT ATG GGC AAC CTT CAA AAG GAC TCT CCA
 E G G D E V L V G F V M G N L Q K D S P
 1321/441
 TTC ATA TTA GTA TTA ATT CAT ATA TAT GCA GCG CCA AAT TCC AGA CAT TGA TAT GCT CTC
 F I L V L I H I Y A A P N S R H * Y A L
 1381/461
 AAA TAG GAT GTT CTG CTC CCC TCT TGT ATT TGT ATG CAG GAA AGG GTT TGT AGG GAG TTT
 K * D V L L P S C I C M Q E R V C R E F
 1441/481
 ACC CCC CCC CCC CCC GCC TTT CTT TGG GGA AGA AAG ACA TAT TCT GGA AGC CTT
 T P P P P P A F L W G R K T Y S G S L
 1501/501
 CCA GTA GTT CAA AA
 P V V Q

FIG. 1 (3/3)

4/8

FIG. 2

plsb	-	Y	F	V	E	G	G	R	S	R	T	G	R	L	L	D	-
	
plsc	-	M	F	P	E	G	T	R	S	R	G	R	G	L	L	P	-
	
maize	-	L	F	V	E	G	T	R	F	T	P	A	K	L	L	A	-
	
		*	*	+	*	*	+	*	+	+				*	*		

5 / 8
FIG. 3A: mutant
JC201B: mutant +
pPLSCC: mutant +
maize
cDNA

6 / 8

FIG. 4




FIG. 5 (1/2)

	190	200	210	220	230	240
maize	LLAAQEYAAASQGLPAPRNVLI	PRTKGFVSAVSIMRDFVPAIYDT	TVIVPKDSPQPTMLRI			
	:	:	:	:	:	:
rape	LKAAQEYAAASSELVPVRNVLI	PRTKGFVSAVSNMRSFVPAIYDM	TVVAIPKTSPPPTMLRL			
	110	120	130	140	150	160
	250	260	270	280	290	
maize	LKGQSSVIHVRMKRRHAMSEMPK	SDDEVSKWCKDIFVAKDALLDKHL	ATGTF-DEEIRPIG			
	:	:	:	:	:	:
rape	FKGQPSVVHVHIKCHSMKDLPE	SEDEIAQWCRDQFVTKDALLDKH	IAADTFAGQKEQNIG			
	170	180	190	200	210	220
	300	310	320	330	340	350
maize	RPVKSLVTLFWSCLLLEFGAIE	FFKWTQLLSTWRGVAF	TAAAGMALVTGVMHVFIMFSQAE			
	:	:	:	:	:	:
rape	RPIKSLAVVLSWACLLTLGAMK	FLHWSNLFSSWKGI	ALSALGLGIITLCMQILIRSSQSE			
	230	240	250	260	270	280
	360	370				
maize	RSSSARAARNRVKKE					
	:	:	:	:	:	:
rape	RSTPAKVAPAKPKDN					
						290

8 / 8

FIG. 5(2/2)

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/GB 93/02528

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C12N15/54 C12N15/82 C12N15/11 C12N9/10 A01H5/00
C11B1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C12N A01H C11B

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
O, X	FAT SCIENCE TECHNOLOGY. 47TH ANNUAL MEETING OF THE GERMAN SOCIETY FOR FAT SCIENCE, 2-5 SEPTEMBER, 1991. vol. 93, no. 11, November 1991 pages 417 - 418 PETEREK, G., ET AL. 'Wege zur Klonierung der 1-Acylglycerin-3-Phosphat-Acyltransferase'	1-4, 14, 15
Y	see the whole document --- -/--	6-13, 16, 17

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* "&" document member of the same patent family

Date of the actual completion of the international search

2 March 1994

Date of mailing of the international search report

27-04-1994

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

Intern: J Application No

PCT/GB 93/02528

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>BIOL. CHEM. HOPPE-SEYLER. WORKSHOP HELD SEPTEMBER 10-13, 1991. vol. 372, no. 8 , 1991 page 539 PETEREK, G., ET AL. 'Approaches of cloning the 1 acylglycerol-3-phosphate acyltransferase' see the whole document ---</p>	6-13, 16, 17
Y	<p>FAT SCIENCE TECHNOLOGY vol. 93, no. 8 , August 1991 pages 288 - 290 WOLTER, F.P., ET AL. 'Biochemical and molecular biological approaches for changing the fatty acid composition of rape seed oil' see the whole document ---</p>	6-13, 16, 17
A	<p>WO,A,92 13082 (KIRIN) 6 August 1992 see the whole document ---</p>	1-4, 6-13, 16, 17
A	<p>CHEMICAL ABSTRACTS, vol. 116, no. 9, 1992, Columbus, Ohio, US; abstract no. 79057, LOEHDEN, I., ET AL. 'Acyl-CoA:1-acylglycerol-3-phosphate acyltransferase from developing seeds of Limnanthes douglasii (R.Br.) and Brassica napus (L.)' see abstract & PLANT LIPID BIOCHEM., STRUCT. UTIL. PROC. INT. SYMP. PLANT LIPIDS 9TH., EDITED BY QUINN, P.J., ET AL. 1990 pages 175 - 177 ---</p>	5
A	<p>CHEMICAL ABSTRACTS, vol. 113, no. 3, 1990, Columbus, Ohio, US; abstract no. 20895, BERNERTH, R., ET AL. 'Utilization of erucoyl-CoA by acyltransferases from developing seeds of Brassica napus (L.) involved in triacylglycerol biosynthesis' see abstract & PLANT SCI. (LIMERICK, IREL.) vol. 67, no. 1 , 1990 pages 21 - 28 ---</p>	5

-/--

INTERNATIONAL SEARCH REPORT

Interns I Application No
PCT/GB 93/02528

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 115, no. 19, 1991, Columbus, Ohio, US; abstract no. 201740, HARES, W., ET AL. 'Substrate specificities of the membrane-bound and partially purified microsomal acyl-CoA:1-acylglycerol-3-phosphate acyltransferase from etiolated shoots of Pisum sativum (1.)' see abstract & PLANTA vol. 185, no. 1 , 1991 pages 124 - 131 ---	5
A	MOLECULAR AND GENERAL GENETICS vol. 232, no. 2 , March 1992 , BERLIN DE pages 295 - 303 COLEMAN, J. 'Charcterization of the Escherichia coli gene for 1-acyl-sn-glycerol-3-phosphate acyltransferase (plsC)' see the whole document ---	1
A	PLANT PHYSIOLOGY vol. 91 , 1989 pages 1288 - 1295 OO, K-C, ET AL. 'Lysophosphatidate acyltransferase activities in the microsomes from palm endosperm, maize scutellum, and rapeseed cotyledon of maturing seeds' see the whole document -----	5

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/GB 93/02528

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9213082	06-08-92	AU-A- 1163392 EP-A- 0567648	27-08-92 03-11-93

This Page is inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images
problems checked, please do not report the
problems to the IFW Image Problem Mailbox**

This Page Blank (uspto)